

## Xpert AmpliFi

(dNTPs included)

#GE17.0100 (100 U) | GE17s (trial size)  
 (FOR RESEARCH ONLY)



**Product:** Xpert AmpliFi DNA Polymerase is an all-round and robust high-fidelity enzyme with enhanced DNA binding, resulting in improved processivity, shorter extension times, higher yield and capacity to amplify more difficult and longer targets. Combined with its extremely low error-rate, this makes Xpert AmpliFi ideal for applications such as high-fidelity PCR, site-directed mutagenesis, cloning, long-range PCR and DNA sequencing, when robustness and proof-reading are extremely important.

Xpert AmpliFi DNA polymerase has been derived from Pfu DNA polymerase for its proof-reading activity. Xpert AmpliFi has a very high accuracy resulting in fidelity that is about 100x higher than regular Taq DNA polymerase. DNA binding has been improved by protein engineering resulting in shorter extension times (2-6kb/min), making this also an ideal choice for consistent results in fast PCR amplifications.

**Applications:** High-Fidelity PCR, cloning, site-directed mutagenesis. Long-Range PCR, DNA sequencing

**Contents:** #GE17.0100 comprises 100U of Xpert AmpliFi DNA Polymerase at a concentration of 2U/μl. The supplied reaction buffer (5x) includes not only MgCl<sub>2</sub>, but also dNTPs, enhancers and stabilizers, optimized to increased PCR success rates. #GE17s is a trial sample (20U).

Component	GE17.0100	GE17s
Xpert AmpliFi (2U/μl)	50μl	10μl
AmpliFi Reaction Buffer (5x)	1.7ml	200μl
AmpliFi Enhancer Mix (10x)	1.7ml	300μl

Xpert AmpliFi is supplied with 1 vial of Reaction Buffer (5x), consisting of optimized concentrations of MgCl<sub>2</sub>, dNTPs, enhancers and stabilizers and 1 vial of Enhancer Mix for difficult and/or long targets (e.g., GC-rich or with complex secondary structures).

**QC:** Functionally tested in PCR. Absence of endonucleases, exonucleases, and ribonucleases was confirmed by appropriate assays

**Properties:**

- Amplicon size: up to 20kb
- Extension rate: 2-6kb/min
- Hotstart: No
- A-overhang: No
- Error-rate: 100x lower than normal Taq

**Storage:** -20°C and protected from light for at least 1 year. No loss of performance is detected after 20 freeze/thaw cycles.

### Prior to use

Optimal PCR cycling conditions (incubation times, temperatures, and concentrations) depend on DNA target (GC-content, size, quantity, purity, etc.), specific primers, dNTPs, buffer composition and MgCl<sub>2</sub>-concentration, and need to be determined case by case. AmpliFi Reaction Buffer already includes dNTPs and has been already been optimized with respect to the MgCl<sub>2</sub> concentration and other components to maximize success rates. Thus, it is not recommended to add additional MgCl<sub>2</sub> or dNTPs. We suggest starting with the following basic protocol and subsequently optimizing annealing temperature, incubation times and cycling number. In case no amplification is observed, it is recommended to add AmpliFi Enhancer Mix as it may improve performance especially for some difficult and/or long templates.

## Basic Protocol


1. Mix for each PCR reaction:

Component	Volume (25µl)	Final Conc.
AmpliFi Reaction Buffer (5x)	5 µl	1X
Forward primer (5 pmol/µl)	2 µl	0.4 µM
Reverse primer (5 pmol/µl)	2 µl	0.4 µM
Template DNA*	0.25 – 10 µl	1-250 ng*
Xpert AmpliFi DNA Polymerase (2U/µl)	0.25 µl	0.05 U/µl
AmpliFi Enhancer Mix (10x) <b>[optional]</b>	2.5 µl	1X
PCR-grade water	up to 25 µl	

\*) In case of gDNA <100ng (total amount) and in case of complex DNA using AmpliFi Enhancer <5ng  
For smaller/larger reaction volumes, scale it down/up proportionally.

In order to minimize risk of contamination, reagent loss and improve pipetting accuracy, we recommend to prepare a mastermix for multiple samples (N), always including a negative control for the detection of possible contaminants, by mixing all components (N+1), except template DNA, dividing the mixture equally into each tube and then add template DNA or PCR grade water in case of the control to the individual PCR tubes.

2. Set-up initial PCR amplification as follows:



After an initial cycle of 1 min at 95°C (denaturation of template DNA including removal of all secondary DNA structures such as hairpins), cycle 25-35 times for 15 seconds at 95°C, 15 seconds at Ta, and 30 seconds per kb of target DNA at 72°C for extension. Set the annealing temperature (Ta) as the melting temperature (Tm) of the primer with the lowest Tm. After amplification, include a final extension step of 3 min at 72°C. Analyze PCR products by DNA Agarose gel electrophoresis.

## Optimization

### Denaturation

If amplification of GC-rich templates results in low yield, increasing temperature to 98°C-100°C may improve amount of product.

### Annealing Temperature (Ta) and Primers

Optimizing the annealing temperature is crucial, as a too low temperature might result in non-specific amplification whereas a too high temperature results in no amplification. The melting temperature (Tm) is defined as the temperature in which 50% of the primer and its complementary sequence of the target DNA are present as duplex DNA. By increasing the temperature above the melting temperature, this percentage decreases, however, primers will still anneal (up to a certain point) and initiate extension. PCR can therefore be performed at temperatures of several degrees higher than Tm and it is therefore recommended to optimize the Ta by performing a temperature gradient (e.g. starting at the lowest Tm and increasing with 2°C increments). Ideally, primers have melting temperatures of approximately 60°C and final concentration should be between 0.2 and 0.6µM (each).

### Incubation times.

Denaturation and annealing steps may require less time depending on the thermocycler apparatus, reaction volume and PCR tube (varies with the efficiency of heat-transfer). It might be worthwhile to optimize (reduce) times to as low as 10 seconds for both denaturation and annealing steps, which will greatly reduce overall PCR time. Optimal extension is carried out with a velocity of 2kb/min but may be reduced to as little as 6kb/min for less demanding templates.